Peertechz



GLOBAL JOURNAL OF Medical and Clinical Case Reports @ SPENALCESS

Case Report

Bone and Joint Infection Caused by Clindamycin and Penicillin-G Resistant, **Toxigenic** Corynebacterium ulcerans

José Bras-Cachinho^{1,2}, Coralie Lemaire^{1,3}*, Laura Chaufour^{1,4}, Nadège Lépine^{1,3}, Louis Bernard^{4,5} and Marie-Frédérique Lartique^{1,3,4}*

¹Department of Bacteriology-Virology-Hospital Hygiene, Tours University Hospital, Tours, France ²Chartres Hospitals, Chartres, Centre-Val de Loire, France

³University of Tours, INRAE, Infectiology and Public Health, BRMF, Tours, France

⁴Tours Study Group Member of CRIOGO (Reference Center for Osteoarticular Infections), Tours, Centre-Val de Loire, France

5CHRU of Tours. Tours. France

Received: 21 February, 2025 Accepted: 03 March, 2025 Published: 04 March 2025

*Corresponding authors: Coralie Lemaire, Department of Bacteriology-Virology-Hospital Hygiene, Tours University Hospitals, Tours, France, E-mail: c.lemaire@chu-tours.fr

ORCiD: https://orcid.org/0009-0007-9810-2695

Marie-Frédérique Lartique, Department of Bacteriology-Virology-Hospital Hygiene, Tours University Hospital, Tours, France, E-mail: mariefrederique.lartigue@univ-tours.fr

ORCiD: https://orcid.org/0000-0003-4948-7959

Keywords: Corynebacterium ulcerans; Bone and joint infection; Tox gene

Copyright License: © 2025 Bras-Cachinho J. et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License. which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

https://www.clinsurggroup.us

Check for updates

Abstract

Corynebacterium diphtheriae, C. ulcerans, and C. tuberculosis are the three species belonging to the diphtheriae complex. C. While C. diphtheriae infections are becoming rare in developed countries, thanks to vaccination, C. ulcerans and C. pseudotuberculosis infections still are rare but increase and are often caused by contact with domestic animals. It should be noted, however, that vaccination protects against diphtheria toxin-producing strains. Here we report an uncommon case of bone and joint infection caused by toxigenic Corynebacterium ulcerans resistant to both clindamycin and penicillin G, in a vaccinated patient. This patient owned a dog whose samples showed carriage of toxigenic C. ulcerans.

Introduction

In developed countries, diphtheria due to C. diphtheriae is now a rare disease, thanks to vaccination. Nevertheless, one should not forget that diphtheria and diphtheria-like diseases can also be caused by other toxigenic corynebacteria: Corynebacterium pseudotuberculosis and Corynebacterium ulcerans. The latter is increasingly reported in industrialized countries as shown in some reviews where the cases listed, with a few exceptions, all come from these countries [1,2]. This can be explained by the lack of diagnostic resources in other countries, but also by the fact that transmission between domestic animals and humans is increasing, or better diagnosed, in developed countries. Various domestic animals are carriers of C.ulcerans, but dogs in particular, as shown by the 2023 activity report and studies

from the French National Corynebacteria Reference Center [3,4]. Other studies have shown the link between these strains isolated from humans and those from their pets, using various comparison techniques such as ribotyping or next-generation sequencing [5,6]. Analysis of antibiotic resistance in strains isolated from dogs, showing resistance to certain antibiotics [7], in particular penicillin and clindamycin, as we describe here with an uncommon case of Bone and Joint Infection (BJI).

Case report

A 74-year-old man was admitted to the Trousseau University Hospital (Tours, France) in January 2021 for the surgical treatment of a left tibiotalar arthrosis, by a Talaris total ankle prosthesis. His medical history includes vitamin K antagonists for atrial fibrillation and two hip and knee joint prostheses.

Three months after the ankle replacement, the patient complained about a swollen anterior scar with suppuration for about three days, which led him to be urgently hospitalized in order to be revised surgically.

During surgery, bacteriological sampling was performed (*n* = 5), and all cultures became positive within 48 hours. Matrixassisted laser desorption/ionisation time-of-flight mass spectrometry (MALDI-TOF, Bruker Daltonics GmbH, Bremen, Germany) identified *Corynebacterium ulcerans*, with an excellent score (>2) using the MALDI Biotyper IVD library. Additional tests were performed by the *Corynebacterium* French National Reference Center (Institut Pasteur – Paris) which confirmed the initial identification and detected the presence of the *tox* gene by qPCR.

A throat swab taken two days later to search for a potential colonization site was negative for *C. ulcerans*. No other sites were tested.

Empirical treatment combined piperacillin/tazobactam and linezolid. Once *C. ulcerans* was identified, an early switch to amoxicillin/clavulanic acid was carried out to treat this microorganism and the other anaerobes present in one of the samples. The strain was susceptible to amoxicillin (MIC = 0.19 mg/L, E-Test Biomérieux, Marcy-l'Étoile, France), ciprofloxacin, vancomycin, trimethoprim-sulfamethoxazole, and linezolid, but resistant to penicillin G (MIC = 0.25 mg/L) and clindamycin (MIC = 2 mg/L).

The patient's condition improved but complete recovery was not achieved. Particularly necrotic tissue appeared on the scar. A second surgery was therefore performed to trim the scar ten days after the first one, with new bacteriological sampling. All the new samples (n = 5) were negative for *C. ulcerans*.

The evolution was favorable, and treatment with amoxicillin 2 g three times a day for three months was pursued.

Further questioning of the patient, a former farmer, revealed that he owns a dog. In addition, the patient received his last diphtheria vaccine booster in January 2021.

In order to substantiate the origin of the infection, the veterinarian was contacted to collect canine samples which, although the dog was asymptomatic, were positive for toxigenic *C. ulcerans.* The dog was also treated with amoxicillin.

C. ulcerans is a fastidious, facultative aero-anaerobic Gram-positive bacillus, first isolated in 1926. *C. ulcerans*, such as *Corynebacterium diphtheriae* and *Corynebacterium pseudotuberculosis*, harbor a lysogenic phage that carries the tox gene, encoding an exotoxin. These three species can be responsible for diphtheria, a potentially lethal disease caused by toxigenic strains. In France, thanks to very high vaccination coverage (98% of children [3]), diphtheria due to *C. diphtheriae* has disappeared, with all 24 infections observed between 2002 and 2018 being imported cases [8]. However, diphtheria-

like infections due to toxigenic *C. ulcerans* have outnumbered those caused by toxigenic *C. diphtheriae* in many industrialized countries [6,9].

C. ulcerans has been isolated from a variety of domestic and wild animals, and although its transmission is mainly zoonotic, human-to-human transmission has been reported [1,10]. Historically, the main sources of *C. ulcerans* infection were consumption of unpasteurized milk or exposure to livestock [11], but pets are increasingly involved in infections [3].

Interestingly, our strain was resistant to both penicillin G and clindamycin. This seems to be rare. According to the annual activity report of the *Corynebacterium* National Reference Center, 13 isolates of toxigenic *C. ulcerans* were confirmed between 2016 and 2018 in France [8]. Although clindamycin resistance seems to be frequent (33 to 60%) [3], only one strain was resistant to both penicillin G and clindamycin. Furthermore, only another strain was also isolated from bone and joint infection (BJI) during this period.

Data suggest that diphtheria vaccination may protect against the development of disease caused by toxigenic *C. ulcerans* [12]. Nevertheless, the patient had received a diphtheria vaccine booster three months before the onset of symptoms. Despite an immune response measured at 3.87 IU/mL (EIA Virotech, Russelsheim, Germany) (threshold = 0.10 UI/mL), he still developed a BJI.

Discussion

This case of BJI infection caused by C. ulcerans is rare as shown in some reviews and articles. Indeed in developed countries, most human infections involving C. ulcerans are respiratory infections, sometimes similar to diphtheria with pseudomembranous angina, or cutaneous and subcutaneous infections that can lead to abscessation [1,13]. Vaccination is directed against diphtheria toxins and is therefore effective against C. ulcerans strains producing these toxins. It is therefore effective against the effects of diphtheria toxin but does not protect against carriage or infection by non-toxigenic strains. Moreover, the fact that a strain carries the tox gene does not mean that the toxin is necessarily produced and secreted by the strain and also that the toxin may have a slightly different molecular structure [14]. Moreover, vaccination can eradicate cases of diphtheria caused by C. diphtheriae, since the carrier is strictly human, but this is not the case for C. ulcerans, whose reservoir is zoonotic, and animals remain the main source of contamination for humans [3,13,15]. The question of vaccination in animals may arise, but the reservoir is very large, and covering all animals to eradicate carriage seems utopian. Perhaps vaccination of companion animals (dogs and cats) could be considered, but vaccination coverage would have to be high to be effective.

Conclusion

In conclusion, we report an uncommon case of BJI caused by toxigenic *C. ulcerans* resistant to both clindamycin and penicillin *G.* Although molecular comparison wasn't possible between the patient's strain and that of his dog, it seems very likely that the patient's dog was the source of the infection.

Citation: Bras-Cachinho J, Lemaire C, Chaufour L, Lépine N, Bernard L, Lartigue MF. Bone and Joint Infection Caused by Clindamycin and Penicillin-G Resistant, Toxigenic Corynebacterium ulcerans. Glob J Medical Clin Case Rep. 2025:12(3):070-072. Available from: https://dx.doi.org/10.17352/2455-5282.000202 This represents a limitation of the study since the source of infection is highly probable but cannot be formally confirmed.

Further clinical and microbiological investigations are required to evaluate the protection conferred by diphtheria vaccination, regarding *C. ulcerans* infections.

Acknowledgment

The authors thank the laboratory technicians who helped to identify this pathogen, the Corynebacterium French National Reference Center (Institut Pasteur – Paris) and the Tours study group of CRIOGO.

Ethical considerations declaration

The patient's consent was obtained to publish this case report.

Author contributions

José Bras-Cachinho: acquisition of data, analysis and interpretation, writing of the manuscript.

Louis Bernard, Laura Chaufour, Coralie Lemaire, and Nadège Lépine: revision of the document, final approval for publication.

Marie-Frédérique Lartigue: writing of the manuscript, revision of the document, final approval for publication.

References

- Hacker E, Antunes CA, Mattos-Guaraldi AL, Burkovski A, Tauch A. Corynebacterium ulcerans, an emerging human pathogen. Future Microbiol. 2016;11:1191–1208. Available from: https://doi.org/10.2217/fmb-2016-0085
- Wagner KS, White JM, Crowcroft NS, De Martin S, Mann G, Efstratiou
 A. Diphtheria in the United Kingdom, 1986-2008: the increasing role of Corynebacterium ulcerans. Epidemiol Infect. 2010;138:1519–1530. Available from: https://doi.org/10.1017/s0950268810001895
- CNR activity reports on Corynebacteria of the diphtheriae complex. Institut Pasteur. 2023. Available from: https://www.pasteur.fr/fr/sante-publique/ CNR/les-cnr/corynebacteries-du-complexe-diphtheriae/rapports-d-activite
- Museux K, Arcari G, Rodrigo G, Hennart M, Badell E, Toubiana J, et al. Corynebacteria of the diphtheriae species complex in companion animals:

clinical and microbiological characterization of 64 cases from France. Microbiol Spectr. 2023;11(3):e0000623. Available from: https://doi.org/10.1128/spectrum.00006-23

- Bonmarin I, Guiso N, Le Flèche-Matéos A, Patey O, Patrick AD, Levy-Bruhl D. Diphtheria: a zoonotic disease in France? Vaccine. 2009;27:4196–4200. Available from: https://doi.org/10.1016/j.vaccine.2009.04.048
- Meinel DM, Margos G, Konrad R, Krebs S, Blum H, Sing A. Next generation sequencing analysis of nine Corynebacterium ulcerans isolates reveals zoonotic transmission and a novel putative diphtheria toxin-encoding pathogenicity island. Genome Med. 2014;6:113. Available from: https://doi.org/10.1186/s13073-014-0113-3
- Katsukawa C, Komiya T, Yamagishi H, Ishii A, Nishino S, Nagahama S, et al. Prevalence of Corynebacterium ulcerans in dogs in Osaka, Japan. J Med Microbiol. 2012;61:266–273. Available from: https://doi.org/10.1099/jmm.0.034868-0
- 8. Diphtérie. Available from: https://www.santepubliquefrance.fr/maladies-ettraumatismes/maladies-a-prevention-vaccinale/diphterie
- Zakikhany K, Efstratiou A. Diphtheria in Europe: current problems and new challenges. Future Microbiol. 2012;7:595–607. Available from: https://doi.org/10.2217/fmb.12.24
- Konrad R, Hörmansdorfer S, Sing A. Possible human-to-human transmission of toxigenic Corynebacterium ulcerans. Clin Microbiol Infect. 2015;21:768– 771. Available from: https://doi.org/10.1016/j.cmi.2015.05.021
- 11. Tiwari TS, Golaz A, Yu DT, Ehresmann KR, Jones TF, Hill HE, et al. Investigations of 2 cases of diphtheria-like illness due to toxigenic Corynebacterium ulcerans. Clin Infect Dis. 2008;46:395–401. Available from: https://doi.org/10.1086/525262
- 12. Otshudiema JO, Acosta AM, Cassiday PK, Hadler SC, Hariri S, Tiwari TSP. Respiratory illness caused by Corynebacterium diphtheriae and C. ulcerans, and use of diphtheria anti-toxin in the United States, 1996-2018. Clin Infect Dis. 2020;ciaa1218. Available from: https://doi.org/10.1093/cid/ciaa1218
- Yamamoto A, Hifumi T, Ato M, Iwaki M, Senoh M, Hatanaka A, et al. Clinical characteristics of Corynebacterium ulcerans infection, Japan. Emerg Infect Dis. 2023;29:1505–1515. Available from: https://doi.org/10.3201/eid2908.220058
- 14. La maladie Recommandations CNR Corynebactérie. Institut Pasteur. 2016. Available from: https://www.pasteur.fr/fr/sante-publique/CNR/les-cnr/ corynebacteries-du-complexe-diphteriae/la-maladie-recommandations
- Trégoat J-J. Health protection, diseases, drug addiction, epidemiology, vaccination, hygiene. Rev Francoph Lab. 2006;53–54.

Discover a bigger Impact and Visibility of your article publication with Peertechz Publications

Highlights

- Signatory publisher of ORCID
- Signatory Publisher of DORA (San Francisco Declaration on Research Assessment)
- Articles archived in worlds' renowned service providers such as Portico, CNKI, AGRIS, TDNet, Base (Bielefeld University Library), CrossRef, Scilit, J-Gate etc.
- Survival states and the second states of the second states and the
- OAI-PMH (Open Archives Initiative Protocol for Metadata Harvesting)
- Dedicated Editorial Board for every journal
- Accurate and rapid peer-review process
- Increased citations of published articles through promotions
- Reduced timeline for article publication

Submit your articles and experience a new surge in publication services https://www.peertechzpublications.org/submission

Peertechz journals wishes everlasting success in your every endeavours.

072